

VDR gene *FokI* polymorphism as a poor prognostic factor for papillary thyroid cancer

Tumor Biology
November 2018: 1–8
© The Author(s) 2018
Article reuse guidelines:
sagepub.com/journals-permissions
DOI: 10.1177/1010428318811766
journals.sagepub.com/home/tub



Selvihan Beysel^{1,2,3} , Nilnur Eyerci⁴, Ferda Alparslan Pinarli⁴,
Mahmut Apaydin¹, Muhammed Kizilgul¹, Mustafa Caliskan¹,
Ozgur Ozcelik¹, Seyfullah Kan¹ and Erman Cakal¹

Abstract

This is the first study to investigate the effect of vitamin D receptor (*VDR*) gene single-nucleotide polymorphisms on the clinicopathologic features of papillary thyroid cancer in Turkey. A total of 165 patients with papillary thyroid cancer and 172 controls were included in this case-control study. *VDR* gene single-nucleotide polymorphisms *FokI* (rs2228570), *BsmI* (rs1544410), *Apal* (rs7975232), and *TaqI* (rs731236) were evaluated using reverse-transcription polymerase chain reaction. *VDR* gene polymorphisms *BsmI*, *Apal*, and *TaqI* did not differ between the papillary thyroid cancer group and control group ($p > 0.05$, each). *BsmI*, *Apal*, and *TaqI* were not associated with papillary thyroid cancer risk. The *VDR* gene *FokI* CT/TT genotype was associated with an increased papillary thyroid cancer risk (CT vs CC: odds ratio = 1.71, 95% confidence interval = 1.15–2.76, $p = 0.028$; TT vs CC: odds ratio = 2.44, 95% confidence interval = 1.29–4.62, $p = 0.005$; CT/TT vs CC: odds ratio = 1.88, 95% confidence interval = 1.20–2.96, $p = 0.006$; CT/CC vs TT: odds ratio = 1.80, 95% confidence interval = 1.05–3.20, $p = 0.041$). *VDR* gene polymorphisms were not in linkage disequilibrium. The *FokI* TT genotype was associated with having T3 and T4, stage III/IV, extra-thyroidal invasion. The *FokI* CT/TT or TT genotype was associated with developing N1 status, multifocality, tumor size ≥ 10 mm, and treatment with radioiodine therapy. Persistence/recurrence did not differ between the *FokI* genotypes. Carriers of the *FokI* T allele were at an increased risk of more advanced tumor-node-metastasis stage, greater tumor size, multifocality, and extra-thyroidal invasion of papillary thyroid cancer compared with the CC genotype. *VDR* gene *FokI* T allele and TT genotype correlated with aggressiveness of papillary thyroid cancer; thus, *FokI* could be useful as a poor prognostic factor to assess the high risk of papillary thyroid cancer.

Keywords

VDR gene, papillary thyroid cancer, *FokI* polymorphism

Date received: 22 November 2017; accepted: 17 October 2018

Introduction

Thyroid cancer is a relatively rare neoplasm that affects an estimated 14.9 people per 100,000. It is the most common endocrine cancer and constituted predominantly of its papillary subtype. Thyroid cancer accounts for <2% of all cancers in males and 1%–1.5% in females with a female-to-male ratio of 3:1 in almost all ethnic groups.^{1,2} Genetic variation contributes to the development of thyroid cancer; however, the pathogenesis of thyroid cancer is not yet known exactly.^{2,3}

¹Department of Endocrinology and Metabolism, Ankara Diskapi Yildirim Beyazit Teaching and Research Hospital, Ankara, Turkey

²Department of Medical Biology, Başkent University, Ankara, Turkey

³Department of Endocrinology and Metabolism, Afyonkarahisar Sağlık Bilimleri University, Afyonkarahisar, Turkey

⁴Department of Genetic Research, Ankara Diskapi Yildirim Beyazit Teaching and Research Hospital, Ankara, Turkey

Corresponding author:

Selvihan Beysel, Department of Endocrinology and Metabolism, Afyonkarahisar Sağlık Bilimleri University, Afyonkarahisar 03200, Turkey.
Email: beyselselvihan@gmail.com



Creative Commons Non Commercial CC BY-NC: This article is distributed under the terms of the Creative Commons

Attribution-NonCommercial 4.0 License (<http://www.creativecommons.org/licenses/by-nc/4.0/>) which permits non-commercial

use, reproduction and distribution of the work without further permission provided the original work is attributed as specified on the SAGE and Open Access pages (<https://us.sagepub.com/en-us/nam/open-access-at-sage>).

Molecular biomarkers have not been identified for the diagnosis and prognosis of thyroid cancer.⁴

In addition to the effect on calcium and bone homeostasis, 1,25-dihydroxyvitamin D₃ (1,25(OH)₂D₃), the biologically active form of vitamin D, has many extra-skeletal effects such as anti-proliferative, anti-apoptotic, and pro-differentiation properties.⁵⁻⁷ Active vitamin D shows its biologic actions by tightly binding to the intracellular vitamin D receptor (VDR), a hormone-regulated transcription factor. VDR is a member of the nuclear hormone receptor family of transacting transcriptional regulatory factors, including the steroid hormone receptors, thyroid hormone receptors, retinoic acid receptors, and retinoid-X receptors. The *VDR* gene is located on chromosome 12q13.1 and is composed of five promoters, eight coding exons, and six untranslated exons.⁵⁻⁸ Its promoter region is able to generate multiple transcripts. During activation, VDR forms a heterodimer with the related retinoid-X receptor and regulates the expression of many target genes because it binds to the vitamin D response elements in the chromatin region. Active vitamin D regulates gene transcription by binding to the VDR. It shows anti-tumoral effect by stimulating differentiation and apoptosis.⁹ It reduces cellular proliferation, inflammation, angiogenesis, and invasion. VDR regulates other metabolic pathways such as immune response and cancer signaling.⁵⁻¹⁰ These are considered as VDR-mediated signaling pathways. Studies have shown that *VDR* gene polymorphisms are found in patients with cancer.¹¹ *BsmI* (A>G, rs1544410), *Apal* (A>C, rs7975232), *TaqI* (T>C, rs731236), and *FokI* (C>T, rs2228570) are human *VDR* gene polymorphisms. *VDR BsmI*, *Apal*, and *FokI* gene polymorphisms have shown to increase the risk of having breast cancer.¹² *VDR FokI* gene polymorphism increases the susceptibility for prostate cancer, *BsmI* polymorphism does so for malignant melanoma, and *TaqI* increases the risk for renal cell carcinoma.¹¹ The *FokI* polymorphism was associated with an increased ovarian cancer risk.¹³ *FokI* T/T genotype was associated with higher progression rate in head and neck squamous cell carcinoma.¹⁴ *VDR BsmI* polymorphism was a risk factor for colorectal and skin cancer in a Caucasian population and the *Apal* polymorphism was a risk factor for basal cell cancer in an Asian population. The *TaqI* polymorphism was a risk factor for oral, breast, and basal cell cancer, and the *FokI* polymorphism was a risk factor for ovarian and skin cancer.¹⁵ However, to date, the association between the *VDR* gene polymorphism and thyroid cancer has only been investigated in two studies. The *VDR Apal* CC and *FokI* TT polymorphisms were associated with decreased follicular thyroid cancer but haplotype Tabf was associated with an increased risk for follicular thyroid cancer.¹⁰ The *VDR* gene polymorphism

was not associated with differentiated thyroid cancer.¹⁶

This study aimed to investigate the relation between the *VDR* gene polymorphism and the clinical features and prognostic significance of patients with papillary thyroid cancer (PTC) in a Turkish population. This is the first study to investigate the effects of *VDR* gene polymorphisms on the clinical features of PTC including cancer staging, outcomes, pathologic findings, and prognostic factors.

Patients and methods

Study population

A total of 165 patients with PTC (mean age: 46.89 ± 13.22 years) and 172 control subjects (mean age: 45.25 ± 4.89 years) were enrolled in this case-controlled study. The subjects with PTC were recruited among patients who were treated at the Department of Endocrinology and Metabolism at Diskapi Teaching and Research Hospital, Ankara, Turkey, between 2011 and 2015. Patients were followed up for a mean of 39.12 ± 8.4 months. Subjects with anaplastic carcinoma, follicular carcinoma, and nodular hyperplasia were excluded. PTC was confirmed by pathology. Controls with thyroid disease, cancer, autoimmune disease, and severe disease or family history of thyroid carcinoma were excluded. The histopathologic features of thyroid carcinoma were assessed according to tumor size (T), tumor-node-metastasis (TNM) cancer staging, uni-/multi-focal tumor, extra-thyroidal invasion, lymph node metastasis (N), distance metastasis (M), and angiolymphatic invasion. The clinical features of subjects with thyroid carcinoma were assessed according to treatment with/without radioactive iodine treatment. TNM cancer staging and outcomes (disease-free and recurrent/persistent) were assessed according to the American Thyroid Association (ATA) guideline.¹⁷ Clinical follow-up was examined in all patients and outcomes were classified according to disease status. Disease-free survival was defined as negative radiologic examination with a serum thyroglobulin value <1 μg/L and undetectable thyroglobulin antibodies (<40 IU/L). Patients without these criteria were defined as having persistent disease. Tumor recurrence was defined as reappearance of tumor after complete cure. Each subject gave written informed consent. The study was approved by the local ethics committee of Diskapi Teaching and Research Hospital (2015/28).

Genotype analysis

Genotyping of *VDR* gene single-nucleotide polymorphisms (SNPs) *FokI* (rs2228570), *BsmI* (rs1544410), *Apal* (rs7975232), and *TaqI* (rs731236) were performed.

Genomic DNA was isolated from collected peripheral blood samples of the subjects using a DNA Isolation Kit (Roche Diagnostics, Indianapolis, IN, USA). *VDR* polymorphisms were evaluated using reverse transcription-polymerase chain reaction (RT-PCR). *VDR* gene SNPs were separately assessed using a fluorescence-based allele-specific PCR assay, KASPar (KBiosciences, Hoddesdon, UK), performed on a Rotor-Gene Q real-time cyler (Qiagen, Hilden, Germany). Allele discrimination was performed using *Rotor-Gene Q software v.2.3.1* (Qiagen, Hilden, Germany). The genotype identification was performed blind without information on the phenotypes of the subjects.

Statistical analysis

Statistical analysis was performed using the SPSS 18.0 (SPSS, Inc.) software. Variables are presented as mean \pm standard deviation (SD) or median (min–max), percentages (%), odds ratios (OR), and 95% confidence intervals (CIs). Normality was tested using the Kolmogorov–Smirnov and Shapiro–Wilk *W* test. SNPs are expressed as allelic frequency (q) or prevalence of genotypes (%). Categorical variables were analyzed using the chi-square test or Fisher's exact test, where appropriate. Student's t-test was used for normally distributed continuous variables or log-transformed variables between two groups. The Hardy–Weinberg equilibrium (HWE) at individual loci was assessed using the chi-square test. Multiple logistic regression analysis and Fisher's exact test were tested using models: dominant (major allele homozygotes vs heterozygotes + minor allele homozygotes), recessive (major allele homozygotes + heterozygotes vs minor allele homozygotes), and codominant (major allele homozygotes vs heterozygote and minor allele homozygotes vs major allele homozygotes). Statistical significance was defined as *p* value < 0.05 . Pairwise linkage disequilibrium (LD) and correlation coefficients (r^2) were analyzed using the HAPLOVIEW program. We made a variable reflecting all possible combinations of *BsmI*–*ApaI*–*TaqI* genotypes for each SNP.

Results

Age (46.89 ± 13.22 vs 45.25 ± 4.89 years) and sex (women, 81.2% vs 86.8%) did not differ between the patients with PTC and controls ($p > 0.05$; Table 1). $25(\text{OH})\text{D}_3$ levels were lower in patients with PTC than in controls (14.41 ± 8.10 vs 17.47 ± 10.32 ng/mL, $p = 0.005$). Four polymorphisms of the *VDR* gene were HWE in the control group. Minor allele frequency and HWE are shown in Table 2. Distributions of *VDR* gene polymorphisms are shown in Table 3. Frequency of

VDR gene *ApaI* rs7975232, *TaqI* rs731236, and *BsmI* rs1544410 polymorphisms did not differ between patients with PTC and controls in a codominant model and dominant model and recessive model ($p > 0.05$, each). *VDR* gene polymorphisms *ApaI*, *TaqI*, and *BsmI* were not associated with PTC. Frequency of *VDR* gene *FokI* rs2228570 increased in PTC patients than in control ($p < 0.05$). Haploview analysis shows that *VDR* gene polymorphisms are not in LD. Compared to control, *FokI* TT genotype increased in patients with PTC in a recessive model (TT vs CT/CC, OR = 1.80, 95% CI = 1.05–3.20, $p = 0.041$) and *FokI* CT/TT genotype increased in patients with PTC in a dominant model (CT/TT vs CC, OR = 1.88, 95% CI = 1.20–2.96, $p = 0.006$). Compared to control, *FokI* CT and *FokI* TT genotypes increased in patients with PTC in a codominant model (CT vs CC, OR = 1.71, 95% CI = 1.15–2.76, $p = 0.028$ and TT vs CC, OR = 2.44, 95% CI = 1.29–4.62, $p = 0.005$).

Association between *VDR* gene *FokI* rs2228570 and clinical features of PTC patients are shown in Table 4. Patients carrying *FokI* TT genotype increased risk for developing T3 and T4 PTC compared to patients carrying *FokI* CC genotype (TT vs CC: OR = 2.71, 95% CI = 1.13–6.53, $p = 0.003$) and C allele and CC genotype (TT vs CT/CC: OR = 3.21, 95% CI = 1.76–6.09, $p = 0.001$, recessive model). The *FokI* T allele and TT genotype increased risk for developing N1 PTC in wild comparison, dominant and recessive model (TT vs CC: OR = 5.16, 95% CI: 1.31–10.42; $p < 0.001$ and CT/TT vs CC: OR = 4.71, 95% CI: 1.25–12.51, $p = 0.008$; and TT vs CT/CC: OR = 5.14, 95% CI: 2.16–10.59; $p < 0.001$, respectively). Patients carrying the *FokI* TT genotype had increased risk for developing stage III/IV PTC compared with patients carrying the CC genotype (TT vs CC: OR = 4.33, 95% CI: 1.07–10.74; $p = 0.029$) and C allele and CC genotype (TT vs CT/CC: OR = 2.67, 95% CI: 1.04–6.83; $p = 0.036$, recessive model). *FokI* T allele and TT genotype increased risk of multifocal tumor in wild, heterozygote comparison, dominant and recessive model (TT vs CC: OR = 5.20, 95% CI = 1.93–11.94; $p = 0.001$; CT vs CC: OR = 2.72, 95% CI = 1.25–5.93, $p = 0.010$; CT/TT vs CC: OR = 2.26, 95% CI = 1.30–3.76, $p = 0.002$; TT vs CT/CC: OR = 2.55, 95% CI = 1.03–5.93, $p = 0.026$; respectively). The *FokI* T allele and TT genotype were more frequent in patients with PTC with tumors ≥ 10 mm in size compared with tumors < 10 mm in size (TT vs CC: OR = 4.09, 95% CI = 1.58–10.52, $p = 0.003$; CT vs CC: OR = 2.38, 95% CI = 1.10–5.15, $p = 0.026$; CT/TT vs CC: OR = 2.58, 95% CI = 1.25–5.33, dominant model). The *FokI* TT genotype was more common in patients with PTC with extra-thyroidal invasion compared with those without extra-thyroidal invasion (TT vs CC: OR = 15.26, 95% CI = 1.90–30.98; TT vs CT/CC:

Table 1. Characteristics of subjects.

	PTC (n = 165)	Control (n = 172)	p
Age (years) ^a	46.89 ± 13.22	45.25 ± 4.89	0.308
Women (%)	81.2	86.6	0.175
Family history of thyroid cancer (%)	4.9	–	
Preoperative 25(OH)D (ng/mL) ^a	14.41 ± 8.10	17.47 ± 10.32	0.005
Age at the diagnosis (years) ^a	44.70 ± 13.55	–	
Type (%)			
Conventional papillary thyroid cancer	63.6		
Follicular variant of papillary thyroid cancer	32.7		
T classification (%)			
Tx	4.8		
T1 and T2	78.2		
T3 and T4	17.0		
N classification (%)			
Nx	4.8		
N0	78.8–78.2		
N1	16.4–17.0		
M classification (%)			
M1	1.3		
M0	98.7		
Clinical stage (%)			
I and II	85.4		
III and IV	14.6		
Multifocal (%)	56.8		
Cancer size ≥ 1 cm (%)	54.1		
Extra-thyroidal invasion (%)	9.6		
Cervical lymph node metastasis (%)	17.2		
Distance metastasis (%)	1.8		
Capsular invasion (%)	14.7		
Angiolymphatic invasion (%)	8.3		
Perineural invasion (%)	1.9		
I131 radiotherapy (%)	86.1		
Time of follow-up (months) ^a	39.12 ± 8.4		
Disease free (%)	89.1		
Recurrent/persistent (%)	7.9		

PTC: papillary thyroid cancer; T: tumor size; N: lymph node metastasis; M: distance metastasis; 25(OH)D: 25-hydroxy-vitamin D.

Bold value represents significant p-value.

^aData are presented as mean ± SD. Percentage was shown as (%).

Table 2. Minor allele frequency and Hardy–Weinberg equilibrium of VDR gene polymorphisms.

	Risk allele	MAF for study sample	p for HWE in control
<i>Apal</i> rs7975232	C	0.41	0.110
<i>TaqI</i> rs731236	C	0.36	0.122
<i>BsmI</i> rs15444410	G	0.39	0.138
<i>FokI</i> rs2228570	T	0.40	0.269

MAF: minor allele frequency; HWE: Hardy–Weinberg equilibrium.

OR = 2.4, 95% CI = 1.20–4.85, $p < 0.001$, recessive model). The *FokI* TT genotype and T allele were more frequent in patients with PTC treated with radioiodine therapy compared with those without radioiodine therapy (TT vs CT: OR = 3.15, 95% CI = 1.20–8.27, $p = 0.016$; TT vs CC: OR = 7.2, 95% CI = 1.71–21.34, $p = 0.002$; CT/TT vs CC: OR = 5.10, 95% CI = 1.99–8.28, dominant model; TT vs CT/CC: OR = 4.34, 95%

CI = 1.22–9.48, $p = 0.007$, recessive model). Patients with recurrent/persistent disease and disease-free status did not differ between the *FokI* genotypes ($p > 0.05$). The VDR gene polymorphisms *Apal*, *BsmI*, and *TaqI* were not related to pathologic features (multifocality, capsule invasion, extra-thyroidal invasion, TNM-staging, tumor size, lymph node metastasis), radioiodine therapy, and outcome.

Table 3. Genotype analysis of VDR gene polymorphisms.

	Control (%)	PTC (%)	OR (95% CI)	p
<i>Apal</i> rs7975232				
Codominant wild-type AA	30.8	32.7		
Heterozygous AC	54.1	57.6	1.00 (0.62–1.61)	0.991
Homozygous CC	15.1	9.7	0.60 (0.29–1.25)	0.173
Dominant (AA/AC + CC)	30.8/69.2	32.7/67.3	0.91 (0.57–1.44)	0.706
Recessive (AA + AC/CC)	84.9/15.1	90.3/9.7	0.63 (0.31–1.17)	0.132
<i>TaqI</i> rs731236				
Codominant wild-type TT	40.1	37.0		
Heterozygous CT	50.6	50.9	1.09 (0.69–1.72)	0.705
Homozygous CC	9.3	12.1	1.41 (0.67–2.97)	0.359
Dominant (TT/CT + CC)	40.1/59.9	37.0/63.0	1.14 (0.73–1.77)	0.553
Recessive (TT + CT/CC)	90.7/9.3	87.9/12.1	1.34 (0.67–2.69)	0.402
<i>BsmI</i> rs1544410				
Codominant wild-type AA	36.0	30.3		
Heterozygous AG	52.3	55.2	1.25 (0.78–2.01)	0.348
Homozygous GG	11.6	14.5	1.48 (0.73–2.99)	0.265
Dominant (AA/AG + GG)	36.0/64.0	30.3/69.7	1.18 (0.82–2.04)	0.263
Recessive (AA + AG/GG)	88.4/11.6	85.5/14.5	1.29 (0.68–2.44)	0.427
<i>FokI</i> rs2228570				
Codominant wild-type CC	43.6	29.1–28.5		
Heterozygous CT	43.0	49.1–49.1	1.71 (1.15–2.76)	0.028
Homozygous TT	13.4	21.8–22.4	2.44 (1.29–4.62)	0.005
Dominant (CC/CT + TT)	43.6/56.4	29.1/70.9	1.88 (1.20–2.96)	0.006
Recessive (CC + CT/TT)	86.6/13.4	78.2/21.8	1.80 (1.05–3.20)	0.041

PTC: papillary thyroid cancer; OR: odds ratio; CI: confidence interval.

Bold values represent significant p-values.

Table 4. Associations between VDR gene *FokI* polymorphism and clinical feature of papillary thyroid cancer.

Variables	Genotype					Dominant model (CT + TT/CC)		Recessive model (CC + CT/TT)	
	CC (wild)	CT	TT	p	p	OR (95% CI)	p	OR (95% CI)	p
	n = 47	n = 81	n = 37	CT/CC	TT/CC				
T classification (n)									
T1 and T2	39	68	22						
T3 and T4	4	11	13	0.458	0.003	2.60 (0.84–7.99)	0.086	3.21 (1.76–6.09)	0.001
N classification (n)									
N0	40	67	22						
N1	2	12	14	0.088	<0.001	4.71 (1.25–12.51)	0.008	5.14 (2.16–10.59)	<0.001
Clinical stage (n)									
I and II	39	68	27						
III and IV	3	11	9	0.250	0.029	2.28 (0.77–4.78)	0.095	2.67 (1.04–6.83)	0.036
Multifocal (n)	15	47	26	0.010	0.001	2.26 (1.30–3.76)	0.002	2.55 (1.03–5.93)	0.026
Cancer size ≥ 1 cm (n)	15	45	25	0.026	0.003	2.58 (1.25–5.33)	0.009	2.18 (0.98–4.84)	0.052
Extra-thyroidal invasion (n)	1	4	10	0.458	0.001	4.46 (0.66–20.1)	0.057	2.4 (1.20–4.85)	<0.001
Angiolymphatic invasion (n)	1	5	7	0.322	0.010	3.81 (0.57–12.53)	0.094	1.73 (0.96–3.15)	0.005
Recurrent/persistent (n)	1	5	7	0.335	0.065	3.81 (0.57–12.56)	0.095	4.74 (1.48–11.15)	0.062
I131 radiotherapy (n)	30	71	35	0.016	0.002	5.10 (1.99–8.28)	0.001	4.34 (1.22–9.48)	0.007

OR: odds ratio; CI: confidence interval; T: tumor size; N: lymph node metastasis.

The number of subjects is shown as (n).

Bold values represent significant p-values.

Discussion

This is the first study to show the effects of *VDR* gene polymorphisms on the clinical features of PTC including TNM cancer staging, outcomes, pathologic findings, and prognostic factors in a Turkish population. We found that the frequency of *VDR* gene *ApaI*, *TaqI*, and *BsmI* polymorphisms did not differ between PTC and controls with no association. The frequency of *VDR* gene *FokI* TT genotype and CT heterozygotes were increased in PTC compared with the controls. *VDR* gene *FokI* was associated with an increased risk of PTC. Haploview analysis showed that *VDR* gene polymorphisms were not in LD. Patients with PTC carrying the *FokI* TT genotype were more likely to have adverse pathologic and prognostic factors including T3 and T4, stage III/IV, and extra-thyroidal invasion. Patients with PTC carrying the *FokI* CT/TT or TT genotype were more likely to develop N1, multifocal tumor, and tumors ≥ 10 mm. *FokI* CT/TT or TT genotype carriers were more likely to be treated with radioiodine therapy. *VDR* gene *FokI* was related to aggressiveness of PTC disease; however, it was not correlated with persistence/recurrence. Thus, *VDR* gene *FokI* might be used as a biomarker for poor clinicopathologic findings.

In humans, vitamin D shows a supportive effect on thyroid tissue. *VDR* gene *BsmI* and *TaqI* polymorphisms were shown to be associated with decreased risk of autoimmune thyroid disease.¹⁸ *VDR* gene *ApaI*, *BsmI*, and *FokI* polymorphisms were associated with Graves' disease in an Asian population, whereas *ApaI*, *BsmI*, *TaqI*, and *FokI* were not associated with Graves' disease in Caucasians.¹⁹ *TaqI* CC and *FokI* TT were increased in Hashimoto's thyroiditis in Turkey.²⁰ Genetic and epigenetic determinants of thyroid cancer may have an effect on $1,25(\text{OH})_2\text{D}_3$ signaling. Development and progression of thyroid cancer depends on impaired $1,25(\text{OH})_2\text{D}_3$ signaling^{21,22} which has been confirmed in clinical studies^{22–26} and thyroid cancer cell lines.²⁷ Local $1,25(\text{OH})_2\text{D}_3$ -VDR signaling is decreased in primary thyroid cancer with local lymph node metastasis, and complete loss of signaling is more significant in anaplastic thyroid cancer with distant metastasis.^{10,22–25} Hence, thyroid cancer progression is characterized by the loss of proteins that play a role in vitamin D signaling or $1,25(\text{OH})_2\text{D}_3$ sensitivity.²⁷ The signaling effect of $1,25(\text{OH})_2\text{D}_3$ -VDR in thyroid cancer occurs via the local anti-tumoral effect of $1,25(\text{OH})_2\text{D}_3$.^{21,22} This anti-tumoral effect occurs either directly (via VDR binding) or indirectly (via the interaction between other critical transcriptional regulators or cell signaling systems).^{21,28,29} Therapeutic use of vitamin D agonists in thyroid cancer requires the presence of VDR. An experimental study showed that *VDR FokI* CC was associated with vitamin D analog

resistance in the treatment of thyroid cancer.³⁰ Active vitamin D has many anti-cancer features including anti-proliferative, anti-apoptotic, pro-differentiating, and anti-inflammatory effects.^{5–10} These effects are mediated by VDR. VDR regulates also metabolic pathways such as the immune response and cancer signaling.^{5–8} VDR is stimulated by p53, a tumor-suppressing gene.¹⁶ An association was found between vitamin D deficiency and thyroid cancer.^{10,31,32}

Epidemiologic studies reported that there was a relationship between *VDR* gene and breast cancer, prostate cancer, lung cancer, colorectal cancer, and other gastrointestinal cancers.^{4–8} However, reliable data on thyroid cancer are available only in in vitro experimental studies.^{10,23,24,27–31} *VDR* gene *ApaI*, *BsmI*, *TaqI*, and *FokI* genotypes were not associated with risk of PTC in a German population.¹⁰ In an Iranian population, frequency of *VDR* gene *ApaI*, *FokI*, *BsmI*, *TaqI*, and *Tru9* did not differ between differentiated thyroid cancer and controls, with no relationship.¹⁶ However, these studies did not investigate the genotype-phenotype correlation such as TNM cancer staging, outcomes, pathologic findings, and prognostic factors. In this study, vitamin D deficiency was found in patients with PTC compared with controls. This study showed that frequency of *VDR* gene *ApaI*, *TaqI*, and *BsmI* did not differ between patients with PTC and controls. *VDR* gene *ApaI*, *TaqI*, and *BsmI* were not associated with a risk for PTC. *VDR* gene *FokI* rs2228570 showed significant differences between patients with PTC and controls: *VDR* gene *FokI* (variant or heterozygotes) compared with wild type (CC) revealed a significant association. *VDR* gene *FokI* TT (TT vs CC: OR = 2.44, 95% CI = 1.29–4.62, $p = 0.005$) and CT heterozygotes (CT vs CC: OR = 1.71, 95% CI = 1.15–2.76, $p = 0.028$) were associated with an increased PTC risk compared with the controls. Our study suggests that *VDR* gene *FokI* may be associated with the development of PTC. Allele frequency distribution showed a significant association of *FokI* variant allele (T) on susceptibility to PTC. These results revealed that VDR gene *FokI* might play a critical role in the etiology of the PTC. *VDR* gene *FokI* TT genotype was associated with having T3 and T4, stage III/IV, and extra-thyroidal invasion of PTC. *FokI* CT/TT or TT genotype was associated with developing N1, multifocality, and tumor size ≥ 10 mm in PTC. Patients carrying *FokI* CT/TT or TT genotype were mostly treated with radioiodine therapy. The persistence/recurrence rate in PTC did not differ between *FokI* genotypes. Although *VDR* gene *FokI* was associated with more advanced TNM stage, greater diameter of tumor, multifocality, and extra-thyroidal invasion, outcomes in follow-up did not differ. *VDR* gene *FokI* might be considered as a risk for poor clinicopathologic features and advanced stage of PTC. *VDR* gene *FokI* might be suggested as a poor prognostic factor of PTC.

However, the molecular mechanism needs to be identified.

BsmI, *ApaI*, and *TaqI* polymorphisms of VDR gene are found in the three primer untranslated region (3'-UTR) and have been shown to be in strong LD.^{33,34} *FokI* polymorphism was reported as an independent marker of VDR gene because it has not been shown to be in LD with any other VDR polymorphisms.³⁴ Haploview analysis showed that VDR gene polymorphisms were not in LD. *ApaI* and *BsmI* polymorphisms of the VDR gene, both in intron 8, are considered as silent SNPs. These polymorphisms do not change the amino acid sequence of the encoded protein, but they might affect gene expression by modulating stability of messenger RNA (mRNA).¹¹ *TaqI* polymorphism is located at codon 352 in exon 9 of the VDR gene. According to the *Taq* restriction site, products are digested into two or three fragments. TT genotype (absence of restriction site) is related to lower active vitamin D3.⁴⁻¹¹ The only locus that has an impact on the structure of VDR protein is the *FokI* polymorphism, which is located on the 5' end region of VDR gene. VDR gene *FokI* polymorphism is functional because it is found in a coding sequence. *FokI* polymorphism is located in the first ATG starting code of VDR protein. *FokI* is involved in thymine to cytosine (T/C) substitution at exon 2, the first translation initiation region is removed, and the transcriptional activity of VDR is changed.^{4-11,16} It alters the ACG codon that is found 10 base pairs upstream from the translation starting codon and leads to generating an additional starting codon. Two different VDR isoforms occur with the transition of allele T to C in ATG. When the initiating translation starts from this alternative site in the thymine variant, it leads to the generation of a longer VDR protein comprising 427 amino acids. The gene is transcribed in normal length if there is a restriction site. Thus, the C/C allele codes the 424-amino acid protein and the T/T allele codes the 427-amino acid protein. The longer VDR protein has low activity in transcription; accordingly, activation is decreased in the target cell.⁴⁻¹¹ Arai et al.³⁵ reported that compared with *FokI* T/T genotype, *FokI* C/C showed 1.7-fold greater function in vitamin D-dependent transcriptional activation of a reporter through the regulation of a vitamin D response element. *FokI* rs2228570 polymorphism is the only VDR gene polymorphism involved in the generation of an altered protein expression.¹¹

In this study, the control population was consistent with HWE. The small sample size, cross-sectional design, and seasonal change in 25(OH)D₃ are limitations of this study.

In conclusion, this study showed that VDR gene *FokI* might contribute to the susceptibility of PTC risk

in the Turkish population. VDR gene *FokI* correlated with aggressiveness of papillary thyroid cancer but not with outcome of disease. *FokI* could be useful as supplementary prognostic factor to assess high risk of papillary thyroid cancer and to identify patients who need more aggressive management and follow-up. Further studies in different populations are needed to confirm these results.

Availability of data and materials

All data are freely available for scientific purpose.

Acknowledgement

We special thanks to Professor Dr Erkan Yurtcu for genetic analysis of data.


Declaration of conflicting interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The author(s) received no financial support for the research, authorship, and/or publication of this article.

ORCID iD

Selvihan Beyse  <https://orcid.org/0000-0001-6963-1503>

References

1. Ukekwe FI, Olusina DB and Okere PCN. Patterns of thyroid cancers in Southeastern Nigeria: a 15 year histopathologic review (2000–2014). *J Clin Diagn Res* 2017; 11(8): EC16–EC19.
2. Figlioli G, Elisei R, Romei C, et al. A comprehensive meta-analysis of case-control association studies to evaluate polymorphisms associated with the risk of differentiated thyroid carcinoma. *Cancer Epidemiol Biomarkers Prev* 2016; 25: 700–713.
3. Kohler A, Chen B, Gemignani F, et al. Genome-wide association study on differentiated thyroid cancer. *J Clin Endocrinol Metab* 2013; 98(10): E1674–E1681.
4. Bai Y-H, Lu H, Hong D, et al. Vitamin D receptor gene polymorphisms and colorectal cancer risk: a systematic meta-analysis. *World J Gastroenterol* 2012; 18(14): 1672–1679.
5. Gnagnarella P, Pasquali E, Serrano D, et al. Vitamin D receptor polymorphism *FokI* and cancer risk: a comprehensive meta-analysis. *Carcinogenesis* 2014; 35(9): 1913–1919.
6. Raimondi S, Pasquali E, Gnagnarella P, et al. *BsmI* polymorphism of vitamin D receptor gene and cancer risk: a comprehensive meta-analysis. *Mutat Res* 2014; 769: 17–34.
7. Serrano D, Gnagnarella P, Raimondi S, et al. Meta-analysis on vitamin D receptor and cancer risk: focus on the role of *TaqI*, *ApaI*, and *Cdx2* polymorphisms. *Eur J Cancer Prev* 2016; 25(1): 85–96.

8. Tuoresmaki P, Vaisanen S, Neme A, et al. Patterns of genome-wide VDR locations. *PLoS ONE* 2014; 9(4): e96105.
9. Tang C, Chen N, Wu M, et al. FokI polymorphism of vitamin D receptor gene contributes to breast cancer susceptibility: a meta-analysis. *Breast Cancer Res Treat* 2009; 117(2): 391–399.
10. Penna-Martinez M, Ramos-Lopez E, Stern J, et al. Vitamin D receptor polymorphisms in differentiated thyroid carcinoma. *Thyroid* 2009; 19(6): 623–628.
11. Kostner K, Denzer N, Muller CS, et al. The relevance of vitamin D receptor (VDR) gene polymorphisms for cancer: a review of the literature. *Anticancer Res* 2009; 29(9): 3511–3536.
12. Iqbal MUN and Khan TA. Association between Vitamin D receptor (Cdx2, FokI, BsmI, ApaI, BglI, TaqI, and Poly (A)) gene polymorphism and breast cancer: a systematic review and meta-analysis. *Tumour Biol*. Epub ahead of print 26 October 2017. DOI: 10.1177/1010428317731280.
13. Liu Y, Li C, Chen P, et al. Polymorphisms in the vitamin D Receptor (VDR) and the risk of ovarian cancer: a meta-analysis. *PLoS ONE* 2013; 8(6): e66716.
14. Hama T, Norizoe C, Suga H, et al. Prognostic significance of vitamin D receptor polymorphisms in head and neck squamous cell carcinoma. *PLoS ONE* 2011; 6(12): e29634.
15. Xu Y, He B, Pan Y, et al. Systematic review and meta-analysis on vitamin D receptor polymorphisms and cancer risk. *Tumour Biol* 2014; 35(5): 4153–4169.
16. Haghpanah V, Ghaffari SH, Rahimpour P, et al. Vitamin D receptor gene polymorphisms in patients with thyroid cancer. *Gene Ther Mol Biol B* 2007; 11: 299–304.
17. Haugen BR, Alexander EK, Bible KC, et al. 2015 American Thyroid Association Management guidelines for adult patients with thyroid nodules and differentiated thyroid cancer: the American Thyroid Association guidelines task force on thyroid nodules and differentiated thyroid cancer. *Thyroid* 2016; 26(1): 1–133.
18. Feng M, Li H, Chen SF, et al. Polymorphisms in the vitamin D receptor gene and risk of autoimmune thyroid diseases: a meta-analysis. *Endocrine* 2013; 43(2): 318–326.
19. Zhou H, Xu C and Gu M. Vitamin D receptor (VDR) gene polymorphisms and Graves' disease: a meta-analysis. *Clin Endocrinol* 2009; 70(6): 938–945.
20. Yazici D, Yavuz D, Tarcin O, et al. Vitamin D receptor gene ApaI, TaqI, FokI and BsmI polymorphisms in a group of Turkish patients with Hashimoto's thyroiditis. *Minerva Endocrinol* 2013; 38(2): 195–201.
21. Clinckspoor I, Verlinden L, Overbergh L, et al. 1,25-dihydroxyvitamin D3 and a superagonistic analog in combination with paclitaxel or suberoylanilide hydroxamic acid have potent antiproliferative effects on anaplastic thyroid cancer. *J Steroid Biochem Mol Biol* 2011; 124(1–2): 1–9.
22. Clinckspoor I, Hauben E, Verlinden L, et al. Altered expression of key players in vitamin D metabolism and signaling in malignant and benign thyroid tumors. *J Histochem Cytochem* 2012; 60(7): 502–511.
23. Balla I, Tobias B, Kosa JP, et al. Vitamin D-neutralizing CYP24A1 expression, oncogenic mutation states and histological findings of human papillary thyroid cancer. *J Endocrinol Invest* 2015; 38(3): 313–321.
24. Izzhakov E, Somjen D, Sharon O, et al. Vitamin D receptor expression is linked to potential markers of human thyroid papillary carcinoma. *J Steroid Biochem Mol Biol* 2016; 159: 26–30.
25. Khadzokou K, Buchwald P, Westin G, et al. 25-hydroxyvitamin D3 1alpha-hydroxylase and vitamin D receptor expression in papillary thyroid carcinoma. *J Histochem Cytochem* 2006; 54(3): 355–361.
26. Schulten HJ, Al-Mansouri Z, Baghallab I, et al. Comparison of microarray expression profiles between follicular variant of papillary thyroid carcinomas and follicular adenomas of the thyroid. *BMC Genomics* 2015; 16(Suppl. 1): S7.
27. Somjen D, Grafi-Cohen M, Posner GH, et al. Vitamin D less-calcemic analog modulates the expression of estrogen receptors, vitamin D receptor and 1 α -hydroxylase 25-hydroxy vitamin D in human thyroid cancer cell lines. *J Steroid Biochem Mol Biol* 2013; 136: 80–82.
28. Liu W, Asa SL, Fantus IG, et al. Vitamin D arrests thyroid carcinoma cell growth and induces p27 dephosphorylation and accumulation through PTEN/akt-dependent and -independent pathways. *Am J Pathol* 2002; 160(2): 511–519.
29. Verlinden L, Eelen G, Beullens I, et al. Characterization of the condensin component Cnap1 and protein kinase Melk as novel E2F target genes down-regulated by 1,25-dihydroxyvitamin D3. *J Biol Chem* 2005; 280(45): 37319–37330.
30. Sharma V, Fretwell D, Crees Z, et al. Thyroid cancer resistance to vitamin D receptor activation is associated with 24-hydroxylase levels but not the ff FokI polymorphism. *Thyroid* 2010; 20: 1103–1111.
31. Penna-Martinez M, Ramos-Lopez E, Stern J, et al. Impaired vitamin D activation and association with CYP24A1 haplotypes in differentiated thyroid carcinoma. *Thyroid* 2012; 22(7): 709–716.
32. Sahin M, Ucan B, Ginis Z, et al. Vitamin D3 levels and insulin resistance in papillary thyroid cancer patients. *Med Oncol* 2013; 30(2): 589.
33. Song GG and Lee YH. Vitamin D receptor FokI, BsmI, ApaI, and TaqI polymorphisms and susceptibility to ovarian cancer: a meta-analysis. *Immunol Invest* 2013; 42(7): 661–672.
34. Lurie G, Wilkens LR, Thompson PJ, et al. Vitamin D receptor rs2228570 polymorphism and invasive ovarian carcinoma risk: pooled analysis in five studies within the ovarian cancer association consortium. *Int J Cancer* 2011; 128(4): 936–943.
35. Arai H, Miyamoto K, Taketani Y, et al. A vitamin D receptor gene polymorphism in the translation initiation codon: effect on protein activity and relation to bone mineral density in Japanese women. *J Bone Miner Res* 1997; 12(6): 915–921.